arXiv:0709.3544v1 [math-ph] 21 Sep 2007

An analytical solution of Monge differential equation

Rafael Torres-Cordoba
Universidad Autonoma de Cd. Juarez.
Av. Del Charro 450 norte, C.P. 32310
Cd. Juarez Chih. Mexico.

February 2, 2008

Abstract

We present the exact solution to the non linear Monge differential equation $\lambda(x,t)\lambda_x(x,t) = \lambda_t(x,t)$. It is widely accepted that the Monge equation is equivalent to the ODE $\ddot{X}= 0$ of free motion for particular conditions. Furthermore, the Monge Type equations are connected with $X = F(\dot{X}, X; t)$, which can be integrated with quadratures [1]. Other asymptotic solutions are discussed, see e.g. [2].

The solution was reached with calculations that depend upon dimensional representation, which is given by (x,t) coordinates. We present this analytical solution to the Monge differential equation as an implicit solution.

1 Introduction

The Monge equation has been cited in several textbooks for approximately 150 years so far [2], [3], and [1]. Despite its wide number of analytical applications (e.g. Monge-Ampere, which is in a more general form than Monge equation itself), it has only been solved to fulfill particular mathematical conditions. The Monge equation currently exists in the following form:

$$\lambda(x,t)\frac{\partial\lambda(x,t)}{\partial x} = \frac{\partial\lambda(x,t)}{\partial t} \quad or \quad \lambda(x,t)\lambda_x(x,t) = \lambda_t(x,t) \tag{1}$$

Examples of particular solutions to this equation for particular mathematical scenarions are exemplified as follows. [2] first reported that the particular solution to the Monge nonlinear differential equation (1) can be given as:

$$\lambda(x,t) = G(x + \lambda(x,t)t), [2] \tag{2}$$

Leznov et al. [1], demonstrated later on that the free motion Monge-type equations are intrinsically related to $X = F(\dot{X}, X; t)$ when ODE $\ddot{X} = 0$. Such function can be integrated through quadratures, which allows the Monge differential equation's solution to be presented as:

$$x - \lambda(x,t)t = f\langle\lambda(x,t)\rangle \tag{3}$$

Two further examples of trivial solutions to the Monge equation are as follows:

$$x + t\lambda(x,t) = 0 \text{ and } \lambda(x,t) = 0 \tag{4}$$

Finally, Fairlie et. al found another solution to this equation in 1993 [3]. This particular solution can be considered to be the extant most-general solution. Nevertheless, it is not completely accurate, and thus the need for more general, accurate forms to the Monge equation.

$$\lambda(x,t) = \frac{a + x}{1 - t} \tag{5}$$

This solution can be reached upon using an arbitrary differentiable initial value F in terms of a Laplace or Fourier expansion as published in [3]:

$$F(x) = c_0 + c_1 e^{\alpha x} + c_2 e^{2\alpha x} + c_3 e^{3\alpha x} + \cdots, \text{ where } F(0) = c_0. \tag{6}$$

$\lambda(x,t)$ remains to be largely unknown despite of its wide number of citations. We find a precise $\lambda(x,t)$ function through the thorough use of

basic differential geometry procedures. The order of a differential equation is given by the maximum number of times the supposed unknown function has been derived.

The Monge differential equation applications are exemplified in the Witham book, in waves theory, and while obtaining the Bateman differential equation. Fairlie et. al., solved the non linear differential equation of the Monge Equation for special cases of their physical applications, mainly in the field of hydrodynamic problems. He also generalized the equation into a system of equations, which he called the Universal Field Equations [3], [4], [5], [6], [7]. The dispersive deformations of the Monge Equation are studied using ideas originating from topological quantum field theory and the deformation quantization programme, [8], etc.. . ..

2 Statement of results

We begin by standardizing notation and terminology. All results presented on this paper are exact in the sense that integration constants depend only upon the initial conditions. Such dependence will usually be pointed out. The function $\lambda(x,t)$ is the unknown function and only depends upon x and t;

$$\lambda(x,t) = \sin\varphi(x,t) \tag{7}$$

Now substituting (7) into (1), the following equation is obtained;

$$\varphi_x(x,t)\sin\varphi(x,t) = \varphi_t(x,t) \ \ or \ \ \sin\varphi(x,t) = \frac{\varphi_t(x,t)}{\varphi_x(x,t)} \tag{8}$$

Considering Figure 1, (8) is derived with respect to t, to obtain the following:

$$\varphi_t(x,t)\cos\varphi(x,t) = \frac{\partial \frac{\varphi_t(x,t)}{\varphi_x(x,t)}}{\partial t} = \left[\frac{\varphi_t(x,t)}{\varphi_x(x,t)}\right]_t \tag{9}$$

Consequently, we have:

$$\cos\varphi(x,t) = \frac{\varphi_x(x,t)\varphi_{tt}(x,t) - \varphi_{tx}(x,t)\varphi_t(x,t)}{-\varphi_t(x,t)\varphi_x^2(x,t)} \tag{10}$$

See figure2.

Now, taking figure 2 into account and comparing it against Figure 1, the following equations can be deduced:

$$\begin{array}{l} \varphi_x(x,t) = -\varphi_x^2(x,t)\varphi_t(x,t) \ \Longrightarrow \\ \varphi_x(x,t) +_x^2 (x,t)\varphi_t(x,t) = 0 \Longrightarrow \\ \varphi_x(x,t)\left[1 + \varphi_x(x,t)\varphi_t(x,t)\right] = 0 \end{array} \tag{11}$$

Must be complete $\varphi_x(x,t) \neq 0$ then $\varphi_x(x,t)\varphi_t(x,t) = -1$ because if $\varphi_x(x,t) = 0$, then $\varphi(x,t) = f(t)$, which will not satisfy the Monge equation (1),

$$\sqrt{\varphi_x^2(x,t) - \varphi_t^2(x,t)} = \varphi_x(x,t)\varphi_{tt}(x,t) - \varphi_{tx}(x,t)\varphi_t(x,t) \tag{12}$$

and

$$\varphi_t(x,t) = \sqrt{\langle \varphi_x^2(x,t)\varphi_t(x,t)\rangle^2 - \langle \varphi_x(x,t)\varphi_{tt}(x,t) - \varphi_{tx}(x,t)\varphi_t(x,t)\rangle^2} \tag{13}$$

Now substituting (12) into (13), and solving for $\varphi_t(x,t)$:

$$\varphi_x^4(x,t)\varphi_t^2(x,t) = \varphi_x^2(x,t) \implies \varphi_x(x,t)\varphi_t(x,t) = \mp 1 \tag{14}$$

One of the solutions does not exist:

$$\varphi_x(x,t)\varphi_t(x,t) = +1 \tag{15}$$

Because $\varphi = \varphi(x,t)$ is a fuction dependant on x and t, anticonmutative properties with respect to the derivative of x and t can be deduced:

$$\frac{\partial \frac{\partial \varphi(x,t)}{\partial t}}{\partial x} = -\frac{\partial \frac{\partial \varphi(x,t)}{\partial x}}{\partial t} \ or \ \varphi_{xt}(x,t) = -\varphi_{tx}(x,t) \tag{16}$$

which can be proven, as shown in Appendix. If $\lambda(x,t) = \sin\varphi(x,t)$, then

$$\lambda_{xt}(x,t) + \lambda_{tx}(x,t) = \left\{\frac{\partial}{\partial x}, \frac{\partial}{\partial t}\right\}\lambda(x,t) = \left\{\frac{\partial}{\partial t}, \frac{\partial}{\partial x}\right\}\lambda(x,t) = -2\lambda(x,t) \tag{17}$$

The correct solution can then be presented as:

$$\varphi_x(x,t)\varphi_t(x,t) = -1 \tag{18}$$

Which complies with equation (11) to deduce the following:

$$\varphi_t(x,t) = -\frac{1}{\varphi_x(x,t)} \tag{19}$$

When $\varphi_{xt}(x,t) = \varphi_{tx}(x,t)$, the commutative properties remain unchanged. Now substituting (19) into (8):

$$\varphi_t(x,t) = i\sqrt{\sin\varphi(x,t)} \text{ and } \varphi_x(x,t) = \frac{-i}{\sqrt{\sin\varphi(x,t)}} \tag{20}$$

By deriving each (20) equation with respect to x and t, respectively, $\varphi_{xt}(x,t) = \frac{\partial\varphi_t(x,t)}{\partial x} = i\frac{\cos\varphi(x,t)}{\sqrt{\sin\varphi(x,t)}}\varphi_x(x,t)$ and $\varphi_{tx}(x,t) = \frac{\partial\varphi_x(x,t)}{\partial t} = -i\frac{\cos\varphi(x,t)}{(\sin\varphi(x,t))^{\frac{3}{2}}}\varphi_t(x,t)$ are obtained. If both equations are brought up together by equaling them to one another, $\varphi_{xt}(x,t) = \varphi_{tx}(x,t) = i\frac{\cos\varphi(x,t)}{\sqrt{\sin\varphi(x,t)}}\varphi_x(x,t) = -i\frac{\cos\varphi(x,t)}{(\sin\varphi(x,t))^{\frac{3}{2}}}\varphi_t(x,t)$, $\varphi_x(x,t) = \frac{\varphi_t(x,t)}{\sin\varphi(x,t)}$ is deduced, which would pretty much be Eq.(8).,

Eq. (20) exactly represents two direct, first order differential equations that are directly integrable:

$$\begin{array}{c} \varphi(x,t) = i\int^t \sqrt{\sin\varphi(x,t')}\; dt' + g(x) \\ \varphi(x,t) = -i\int^x \frac{dx'}{\sqrt{\sin\varphi(x',t)}} + f(t) \end{array} \quad or \tag{21}$$

Because both Eq. (21) solutions are related (i.e. through the equation $\varphi_x(x,t)\varphi_t(x,t) = -1$) by the first derived function with respect to x and t, respectively, this means the solution to the Monge equation must be unique. (21) solution can be seen as follows:

$$\begin{array}{l} t + g(x) = \int^{\varphi(x,t)} \frac{d\varphi'(x,t)}{i\sqrt{\sin\varphi'(x,t)}} = -2\sqrt{2}i\mathbf{F}\left\langle\alpha(x,t), \frac{1}{\sqrt{2}}\right\rangle \quad \text{where} \\ \alpha(x,t) = \arcsin\sqrt{\frac{2\sin\varphi(x,t)}{1+\sin\varphi(x,t)+\cos\varphi(x,t)}} = \arcsin\sqrt{\frac{2\lambda(x,t)}{1+\lambda(x,t)+\sqrt{1-\lambda^2(x,t)}}} \end{array} \tag{22}$$

or

$$\begin{array}{l} x + f(t) = \int^{\varphi(x,t)} i\sqrt{\sin\varphi'(x,t)}d\varphi'(x,t) = -2\sqrt{2}i\mathbf{E}\left\langle\alpha_1(x,t), \frac{1}{\sqrt{2}}\right\rangle + \sqrt{2}i\mathbf{F}\left\langle\alpha_1(x,t), \frac{1}{\sqrt{2}}\right\rangle \quad \text{where} \\ \alpha_1(x,t) = \arcsin\left\langle\sqrt{2}\sin\frac{1}{2}(\frac{\pi}{2} - \varphi(x,t))\right\rangle = \arcsin\left\langle\sqrt{2}\sin\frac{1}{2}(\frac{\pi}{2} - \arcsin\lambda(x,t))\right\rangle \end{array} \tag{23}$$

see e.g. [9], where both solutions given by equations (22) and (23) are equivalent in $\lambda(x,t)$. $g(x)$ and $f(t)$ depend only on the initial conditions, and cannot be zero or constant, because $\lambda(x,t)\lambda_x(x,t) = \lambda_t(x,t)$ must be satisfied. Solutions (22) and (23) are implicit to $\lambda(x,t)\lambda_x(x,t) = \lambda_t(x,t)$. Finally, $\mathbf{F}(\boldsymbol{\beta},\mathbf{a})$ and $\mathbf{E}(\boldsymbol{\beta},\mathbf{a})$ are the Elliptic integrals of the first and second kind, respectively.

References

[1] A. N. Leznov, The connection of Moge-Bateman equations with ordinary differential equations and their generalization, Arxiv: math-phys/9908013 v1, (1999).

[2] Witham G. B. 1974 Linear and Nonlinear Waves, Wiley

[3] David B. Fairlie, Equations of Hydrodynamic type, arXiv:hep-th/9305049 v1 (may 1993).

[4] David B. Fairlie, Implicit solutions to some Lorentz Invariant nonlinear equations revisited, JNMP **12**, 449-456 (2005).

[5] David B. Fairlie, A. N. Leznov, General solution of the Universal Equation in n-Dimensional Space, JNMP **1**, 333-339 (1994).

[6] Thomas Curtright, David Fairlie, Morphing quantum mechanics and fluid dynamics, J. Phys. A Math. Gen **36** 8885-8901 (2003).

[7] Thomas Curtright, David Fairlie, Extra dimensions and nonlinear equations, J, Math. Phys. **44** 2692-2703 (2003).

[8] I.A.B. Strachan, Deformations of the Monge/Riemann hierarchy and approximately integrable systems, arXiv:nlin.SI/0205051 v1 (may 2002).

[9] I.S.Gradshteyn, I.M.Ryzbik, Table of integrals, series, and products ISBN 0-12-294757-6

2.1

appendix

In order to prove $\varphi_{xt}(x,t) = -\varphi_{tx}(x,t)$, using the relation $\varphi_x(x,t)\varphi_t(x,t) = 1$, $\varphi_x(x,t) = \varphi_x^2(x,t)\varphi_t(x,t)$ must be fullfilled. Equation (10) must be modified as it is indicated:

$$\cos\varphi(x,t) = \frac{\varphi_x(x,t)\varphi_{tt}(x,t) - \varphi_{tx}(x,t)\varphi_t(x,t)}{-\varphi_t(x,t)\varphi_x^2(x,t)} = \frac{\varphi_{tx}(x,t)\varphi_t(x,t) - \varphi_x(x,t)\varphi_{tt}(x,t)}{\varphi_t(x,t)\varphi_x^2(x,t)} \tag{24}$$

see figure 3. Now taking into account Figure 3 and Figure1, we obtain:

$$\sqrt{\varphi_x^2(x,t)-\varphi_t^2(x,t)}=\varphi_{tx}(x,t)\varphi_t(x,t)-\varphi_x(x,t)\varphi_{tt}(x,t) \tag{25}$$

from $\varphi_x(x,t)\varphi_t(x,t)=+1$ and Eq. (8), it is found that:

$$\varphi_x(x,t)\varphi_t(x,t)=1 \text{ then } \varphi_t(x,t)=\sqrt{\sin\varphi(x,t)}\text{and } \varphi_x(x,t)=\frac{1}{\sqrt{\sin\varphi(x,t)}} \tag{26}$$

Now substituting (26) into (25):

$$\begin{aligned}
&\sqrt{\varphi_x^2(x,t)-\varphi_t^2(x,t)}=\sqrt{\tfrac{1}{\sin\varphi(x,t)}-\sin\varphi(x,t)}=\sqrt{\tfrac{1-\sin^2\varphi(x,t)}{\sin\varphi(x,t)}}=\\
&=\sqrt{\tfrac{\cos^2\varphi(x,t)}{\sin\varphi(x,t)}}=\tfrac{\cos\varphi(x,t)}{\sqrt{\sin\varphi(x,t)}} \text{ now}\\
&\varphi_{tx}(x,t)\varphi_t(x,t)-\varphi_x(x,t)\varphi_{tt}(x,t)=\tfrac{\cos\varphi(x,t)}{2\sqrt{\sin\varphi(x,t)}}-\tfrac{-\cos\varphi(x,t)}{2\sin\varphi(x,t)}\sqrt{\sin\varphi(x,t)}=\\
&=\tfrac{\cos\varphi(x,t)}{\sqrt{\sin\varphi(x,t)}} \text{ is complete (25), where}\\
&\varphi_{tt}(x,t)=\tfrac{\cos\varphi(x,t)}{2\sqrt{\sin\varphi(x,t)}}\varphi_t(x,t)=\tfrac{\cos\varphi(x,t)}{2} \text{ and}\\
&\ \varphi_{tx}(x,t)=\tfrac{\partial\varphi_x(x,t)}{\partial t}=\tfrac{\partial\frac{1}{\sqrt{\sin\varphi(x,t)}}}{\partial t}=\\
&-\tfrac{\cos\varphi(x,t)}{2\langle\sin\varphi(x,t)\rangle^{\frac{3}{2}}}\varphi_t(x,t)=-\tfrac{\cos\varphi(x,t)}{2\sin\varphi(x,t)}
\end{aligned} \tag{27}$$

From figure 3 and figure 1, we found:

$$\varphi_t(x,t)=\sqrt{\langle\varphi_x^2(x,t)\varphi_t(x,t)\rangle^2-\langle\varphi_x(x,t)\varphi_{tt}(x,t)-\varphi_{tx}(x,t)\varphi_t(x,t)\rangle^2} \tag{28}$$

Now substituting (26) into (28) and using the results of (27):

$$\begin{aligned}
&\varphi_t(x,t)=\sqrt{\sin\varphi(x,t)}\ =\sqrt{\langle\varphi_x^2(x,t)\varphi_t(x,t)\rangle^2-\langle\varphi_x(x,t)\varphi_{tt}(x,t)-\varphi_{xt}(x,t)\varphi_t(x,t)\rangle^2}\\
&=\sqrt{\tfrac{\sin\varphi(x,t)}{\sin^2\varphi(x,t)}-\tfrac{\cos^2\varphi(x,t)}{\sin\varphi(x,t)}}=\sqrt{\tfrac{1-\cos^2\varphi(x,t)}{\sin\varphi(x,t)}}=\sqrt{\sin\varphi(x,t)} \text{ is complete}
\end{aligned} \tag{29}$$

Now we found $\varphi_{xt}(x,t)$ and proved that $\varphi_{xt}(x,t)=-\varphi_{tx}(x,t)$ is as follows:

$\varphi_{xt}(x,t) = \frac{\partial\varphi_t(x,t)}{\partial x} = \frac{\partial\sqrt{\sin\varphi(x,t)}}{\partial x} = \frac{\cos\varphi(x,t)}{2\sqrt{\sin\varphi(x,t)}}\varphi_x(x,t) =$
$= \frac{\cos\varphi(x,t)}{2\sqrt{\sin\varphi(x,t)}}\frac{1}{\sqrt{\sin\varphi(x,t)}} = \frac{\cos\varphi(x,t)}{2\sin\varphi(x,t)}$ but (30)
$\varphi_{tx}(x,t) = -\frac{\cos\varphi(x,t)}{2\sin\varphi(x,t)}$ containen in (27) then
$\varphi_{xt}(x,t) = -\varphi_{tx}(x,t)$ is proved

Now we prove that the second mixed partial derivative of $\lambda(x,t)$ is not independent from the order of differentiation for this case (i.e. that $\lambda_{xt}(x,t) + \lambda_{tx}(x,t) = -2\lambda(x,t)$). From (7), we have

$\lambda(x,t) = \sin\varphi(x,t)$ then $\lambda_x(x,t) = \varphi_x \cos\varphi(x,t)$ and $\lambda_t(x,t) = \varphi_t \cos\varphi(x,t)$ next
$\lambda_{xt}(x,t) = -\varphi_x\varphi_t \sin\varphi(x,t) + \varphi_{xt}\cos\varphi(x,t)$ and $\lambda_{tx}(x,t) = -\varphi_t\varphi_x \sin\varphi(x,t) + \varphi_{tx}\cos\varphi(x,t)$
taking the sum of $\lambda_{xt}(x,t)$ and $\lambda_{tx}(x,t)$ it is obtained
$\lambda_{xt}(x,t) + \lambda_{tx}(x,t) = -\varphi_x\varphi_t \sin\varphi(x,t) - \varphi_t\varphi_x \sin\varphi(x,t)$
where we used $\varphi_{tx} = -\ \varphi_{xt}$ and we know that $\varphi_x\varphi_t = \varphi_x\varphi_t = 1$ then
$\lambda_{xt}(x,t) + \lambda_{tx}(x,t) = -2\sin\varphi(x,t)$ now we using (7), finally $\lambda_{xt}(x,t) + \lambda_{tx}(x,t) = -2\lambda(x,t)$ (31)

This is the reason why equation (15), is not a solution to the Monge Equation (1).

.1 figures



Figure 1: Figure 1


$\sqrt{(\varphi_t\varphi_x^2)^2-(\varphi_x\varphi_{tt}-\varphi_{xt}\varphi_t)^2}$
$-\varphi_t\varphi_x^2$
φ
$\varphi_x\varphi_{tt}-\varphi_{xt}\varphi_t$

Figure 2: Figure 2


$\sqrt{(\varphi_t\varphi_x^2)^2-(\varphi_{tx}\varphi_t-\varphi_x\varphi_{tt})^2}$
$\varphi_t\varphi_x^2$
φ
$\varphi_{tx}\varphi_t-\varphi_x\varphi_{tt}$

Figure 3: Figure 3